

Properties of moments of random variables *

Jean-Marie Dufour [†]
McGill University

First version: May 1995

Revised: April 1999, June 2007, June 2011, May 2016, January 2017

This version: January 2017

Compiled: January 11, 2021, 15:34

* This work was supported by the William Dow Chair in Political Economy (McGill University), the Bank of Canada (Research Fellowship), the Toulouse School of Economics (Pierre-de-Fermat Chair of excellence), the Universitat Carlos III de Madrid (Banco Santander de Madrid Chair of excellence), a Guggenheim Fellowship, a Konrad-Adenauer Fellowship (Alexander-von-Humboldt Foundation, Germany), the Canadian Network of Centres of Excellence [program on *Mathematics of Information Technology and Complex Systems* (MITACS)], the Natural Sciences and Engineering Research Council of Canada, the Social Sciences and Humanities Research Council of Canada, and the Fonds de recherche sur la société et la culture (Québec).

[†] William Dow Professor of Economics, McGill University, Centre interuniversitaire de recherche en analyse des organisations (CIRANO), and Centre interuniversitaire de recherche en économie quantitative (CIREQ). Mailing address: Department of Economics, McGill University, Leacock Building, Room 414, 855 Sherbrooke Street West, Montréal, Québec H3A 2T7, Canada. TEL: (1) 514 398 6071; FAX: (1) 514 398 4800; e-mail: jean-marie.dufour@mcgill.ca. Web page: <http://www.jeanmariedufour.com>

Contents

List of Definitions, Assumptions, Propositions and Theorems

	iii
1. Existence of moments	1
2. Moment inequalities	1
3. Moment-based bounds on tail probabilities	3
3.1. Markov-type inequalities	3
3.2. Markov inequality and ecological fallacies	4
3.3. Markov-type inequalities for bounded variables	5
3.4. Chebyshev-type inequalities	8
3.5. Generalized Markov-type inequalities	10
4. Moments as integrals of distribution and quantile functions	12
5. Integration by parts	15
5.1. Standard results	15
5.2. Extensions	16
6. Tail decay rates and the existence of moments	20
7. Moments of sums of random variables	22
8. Proofs and references	26

List of Definitions, Assumptions, Propositions and Theorems

1.1 Proposition : Existence of absolute and ordinary moments	1
1.5 Proposition : Monotonicity of L_r	1
2.1 Proposition : c_r -inequality	1
2.2 Proposition : Mean form of c_r -inequality	1
2.3 Proposition : Closure of L_r	1
2.4 Proposition : Hölder inequality	2

2.5	Proposition : Cauchy-Schwarz inequality	2
2.6	Proposition : Minkowski inequality	2
2.7	Proposition : Moment monotonicity	2
2.8	Theorem : Liapunov theorem	2
2.9	Proposition : Lower bounds on the moments of a sum	2
2.10	Proposition : Jensen inequality	2
2.11	Proposition : Concave Jensen inequality	2
3.1	Proposition : Markov inequality	3
3.2	Proposition : Tightness of Markov inequality	4
3.3	Proposition : Irrelevance of the mean for quantiles	4
3.4	Proposition : Two-sided Markov-type inequalities for bounded variables	5
3.5	Proposition : Two-sided monotonic Markov-type inequalities	8
3.6	Proposition : Chebyshev inequality	8
3.7	Proposition : Two-sided bounded monotonic Markov-type inequalities	9
3.8	Proposition : Chebyshev inequalities	10
3.9	Theorem : Generic Markov-type inequalities	10
3.10	Proposition : Markov-type inequalities based on truncated moments	11
4.1	Proposition : Distribution decomposition of the first moment	12
4.2	Corollary : Distribution decomposition of the first absolute moment	12
4.3	Proposition : Distribution decomposition of the expected value of $g(X)$	12
4.4	Corollary : Distribution decomposition of r -moments	14
4.5	Corollary : Moment-tail area inequalities	14
4.6	Proposition : Mean-tail area inequalities	14
4.7	Proposition : Integral of the probability of an interval	15
4.8	Corollary : Integral of the probability of a general interval	15
4.9	Proposition : Quantile representation of the mean	15
5.1	Theorem : Existence of Riemann-Stieltjes integral	15
5.2	Theorem : Representation of Riemann-Stieltjes integral for continuous-BV functions	15
5.3	Theorem : Integration by parts	16
5.4	Lemma : Riemann-Stieltjes integration by parts	16
5.5	Lemma : Centered Riemann-Stieltjes integration by parts	16
5.6	Lemma : Improper Riemann-Stieltjes integration by parts	17
5.7	Lemma : Improper centered Riemann-Stieltjes integration by parts	18
5.8	Lemma : Bounded monotonicity condition for tail convergence of an integrable function	19

6.1	Proposition : Tail decay rates based on truncated moments	21
6.2	Corollary : Moment existence and tail area decay	22
6.3	Proposition : Necessary and sufficient condition for the existence of the mean	22
6.4	Proposition : Necessary and sufficient condition for the existence of r -moments	22
7.1	Proposition : Bounds on the absolute moments of a sum of random variables	23
7.2	Proposition : Minkowski inequality for n variables	23
7.3	Proposition : Bounds on the absolute moments of a sum of random variables under conditional symmetry	23
7.4	Proposition : Bounds on the absolute moments of a sum of random variables under martingale condition	24
7.5	Proposition : Bounds on the absolute moments of a sum of random variables under two-sided martingale condition	24
7.6	Proposition : Bounds on the absolute moments of a sum of independent random variables	24
	Proof of Theorem 3.10	26
	Proof of Proposition 4.1	27
	Proof of Theorem 4.3	29
	Proof of Proposition 4.7	31
	Proof of Lemma 5.4	32
	Proof of Lemma 5.5	32
	Proof of Lemma 5.8	32
	Proof of Proposition 7.2	33

Let X and Y be real random variables, and let r and s be real positive constants ($r > 0$, $s > 0$). The distribution functions of X and Y are denoted $F_X(x) = \mathbb{P}[X \leq x]$ and $F_Y(x) = \mathbb{P}[Y \leq x]$.

1. Existence of moments

1.1 EXISTENCE OF ABSOLUTE AND ORDINARY MOMENTS. $\mathbb{E}(|X|)$ always exists in the extended real numbers $\bar{\mathbb{R}} \equiv \mathbb{R} \cup \{\infty\} \cup \{-\infty\}$ and $\mathbb{E}(|X|) \in [0, \infty]$; i.e., either $\mathbb{E}(|X|)$ is a non-negative real number or $\mathbb{E}(|X|) = \infty$.

1.2 $\mathbb{E}(X)$ exists and is finite $\Leftrightarrow \mathbb{E}(|X|) < \infty$.

1.3 $\mathbb{E}(|X|) < \infty \Rightarrow |\mathbb{E}(X)| \leq \mathbb{E}(|X|) < \infty$.

1.4 If $0 < r \leq s$, then

$$\mathbb{E}(|X|^r) < \infty \Rightarrow \mathbb{E}(|X|^s) < \infty. \quad (1.1)$$

1.5 MONOTONICITY OF L_r . $L_s \subseteq L_r$ for $0 < r \leq s$.

1.6 $\mathbb{E}(|X|^r) < \infty \Rightarrow \mathbb{E}(X^k)$ exists and is finite for all integers k such that $0 < k \leq r$.

2. Moment inequalities

2.1 c_r -INEQUALITY.

$$\mathbb{E}(|X + Y|^r) \leq c_r [\mathbb{E}(|X|^r) + \mathbb{E}(|Y|^r)] \quad (2.1)$$

where

$$c_r = \begin{cases} 1, & \text{if } 0 < r \leq 1, \\ 2^{r-1}, & \text{if } r > 1. \end{cases} \quad (2.2)$$

2.2 MEAN FORM OF c_r -INEQUALITY.

$$\begin{aligned} \mathbb{E}\left(\frac{1}{2}(X + Y)^r\right) &\leq \left(\frac{1}{2}\right)^r [\mathbb{E}(|X|^r) + \mathbb{E}(|Y|^r)], & \text{if } 0 < r \leq 1, \\ &\leq \frac{1}{2} [\mathbb{E}(|X|^r) + \mathbb{E}(|Y|^r)], & \text{if } r > 1. \end{aligned} \quad (2.3)$$

2.3 CLOSURE OF L_r . Let a and b be real numbers. Then

$$X \in L_r \text{ and } Y \in L_r \Rightarrow aX + bY \in L_r. \quad (2.4)$$

2.4 HÖLDER INEQUALITY. If $r > 1$ and $\frac{1}{r} + \frac{1}{s} = 1$, then

$$\mathbb{E}(|XY|) \leq [\mathbb{E}(|X|^r)]^{1/r} [\mathbb{E}(|Y|^s)]^{1/s}. \quad (2.5)$$

2.5 CAUCHY-SCHWARZ INEQUALITY.

$$\mathbb{E}(|XY|) \leq [\mathbb{E}(X^2)]^{1/2} [\mathbb{E}(Y^2)]^{1/2}. \quad (2.6)$$

2.6 MINKOWSKI INEQUALITY. If $r \geq 1$, then

$$\mathbb{E}(|X + Y|^r)^{1/r} \leq [\mathbb{E}(|X|^r)]^{1/r} + [\mathbb{E}(|Y|^r)]^{1/r}. \quad (2.7)$$

2.7 MOMENT MONOTONICITY. $[\mathbb{E}(|X|^r)]^{1/r}$ is a non-decreasing function of r , i.e.

$$0 < r \leq s \Rightarrow [\mathbb{E}(|X|^r)]^{1/r} \leq [\mathbb{E}(|X|^s)]^{1/s}. \quad (2.8)$$

2.8 Theorem LIAPUNOV THEOREM. $\log[\mathbb{E}(|X|^r)]$ is a convex function of r , i.e. for any $\lambda \in [0, 1]$,

$$\log[\mathbb{E}(|X|^{\lambda r + (1-\lambda)s})] \leq \lambda \log[\mathbb{E}(|X|^r)] + (1-\lambda) \log[\mathbb{E}(|X|^s)]. \quad (2.9)$$

2.9 LOWER BOUNDS ON THE MOMENTS OF A SUM. If $\mathbb{E}(|X|^r) < \infty$, $\mathbb{E}(|Y|^r) < \infty$ and $\mathbb{E}(Y|X) = 0$, then

$$\mathbb{E}(|X + Y|^r) \geq \mathbb{E}(|X|^r), \quad \text{for } r \geq 1. \quad (2.10)$$

2.10 JENSEN INEQUALITY. If $g(x)$ is a convex function on \mathbb{R} and $\mathbb{E}(|X|) < \infty$, then, for any constant $c \in \mathbb{R}$,

$$g(c) \leq \mathbb{E}[g(X - EX + c)] \quad (2.11)$$

and, in particular,

$$g(EX) \leq \mathbb{E}[g(X)]. \quad (2.12)$$

2.11 CONCAVE JENSEN INEQUALITY. If $g(x)$ is a concave function on \mathbb{R} and $\mathbb{E}(|X|) < \infty$, then, for any constant $c \in \mathbb{R}$,

$$g(c) \geq \mathbb{E}[g(X - EX + c)] \quad (2.13)$$

and, in particular,

$$g(EX) \geq \mathbb{E}[g(X)]. \quad (2.14)$$

3. Moment-based bounds on tail probabilities

3.1. Markov-type inequalities

3.1 Theorem MARKOV INEQUALITY. *Let X be a real random variable such that*

$$\mathbb{P}[X \geq 0] = 1. \quad (3.1)$$

If X has a finite expected value $\mathbb{E}[X]$ and $a > 0$, then

$$\mathbb{P}[X \geq a] \leq \frac{\mathbb{E}(X)}{a}. \quad (3.2)$$

Note inequality (3.2) remains formally valid for $a = 0$ or $\mathbb{E}(X) = 0$, if we adopt the conventions that $0/0 \equiv 1$ and $x/0 = +\infty$ for $x > 0$. It is easy to see that (3.2) entails:

$$\mathbb{P}[X > a] \leq \frac{\mathbb{E}(X)}{a}, \text{ for } a > 0, \quad (3.3)$$

$$\mathbb{P}[X < a] \leq \mathbb{P}[X \leq a] \geq 1 - \frac{\mathbb{E}(X)}{a} = \frac{a - \mathbb{E}(X)}{a}, \text{ for } a \geq \mathbb{E}(X), \quad (3.4)$$

$$\mathbb{P}[X = a] \leq \frac{\mathbb{E}(X)}{a}. \quad (3.5)$$

Further, if $\mathbb{E}(X) > 0$, we can replace a by $a\mathbb{E}[X]$, hence

$$\mathbb{P}[X > a\mathbb{E}(X)] \leq \mathbb{P}[X \geq a\mathbb{E}(X)] \leq \frac{1}{a}, \text{ for any } a > 0, \quad (3.6)$$

$$\mathbb{P}[X < a\mathbb{E}(X)] \leq \mathbb{P}[X \leq a\mathbb{E}(X)] \geq 1 - \frac{1}{a} = \frac{a-1}{a}, \text{ for } a \geq 1, \quad (3.7)$$

$$\mathbb{P}[X = a\mathbb{E}(X)] \leq \frac{1}{a}, \text{ for any } a > 0. \quad (3.8)$$

In general, if we define

$$F_X(x) := \mathbb{P}[X \leq x], \quad (3.9)$$

$$F_X^{-1}(q) := \inf\{x : F_X(x) \geq q\}, \quad 0 < q < 1, \quad (3.10)$$

the Markov inequality entails: for $0 < q < 1$,

$$1 - q \leq \mathbb{P}[X \geq F_X^{-1}(q)] \leq \frac{\mathbb{E}(X)}{F_X^{-1}(q)} \quad (3.11)$$

hence

$$\mathbb{E}(X) \geq (1-q)F_X^{-1}(q) \quad (3.12)$$

or, equivalently,

$$\mathbb{E}(X) \geq qF_X^{-1}(1-q) \quad (3.13)$$

For example, the mean of a positive random variable is at least as large as half its median:

$$\mathbb{E}(X) \geq \frac{1}{2}F_X^{-1}(1/2). \quad (3.14)$$

3.2. Markov inequality and ecological fallacies

Question – Under the assumptions of Theorem 3.1, can we give lower bounds on $\mathbb{P}[X \geq a]$? Clearly, if $\mathbb{E}[X] > 0$,

$$\mathbb{P}[X \geq \mathbb{E}(X)] > 0, \quad (3.15)$$

$$\mathbb{P}[X > a] > 0 \text{ for } 0 \leq a < \mathbb{E}(X). \quad (3.16)$$

This raises the question whether a more informative bound is possible. The answer to this question is no: for any $\mu > 0$ and $0 \leq a < \mu$ and $0 < p < 1$, we can find a nonnegative random variable X such that

$$\mathbb{E}[X] = \mu \text{ and } \mathbb{P}[X \leq a] = p. \quad (3.17)$$

3.2 Proposition TIGHTNESS OF MARKOV INEQUALITY. For any real numbers μ and a satisfying $0 \leq \mu < a$, there exists a nonnegative random variable X such that

$$\mathbb{E}[X] = \mu \text{ and } \mathbb{P}[X \geq a] = \frac{\mu}{a}. \quad (3.18)$$

3.3 Proposition IRRELEVANCE OF THE MEAN FOR QUANTILES. Let X be a random variable such that

$$\mathbb{P}[X \geq 0] = 1. \quad (3.19)$$

If X has a finite expected value $\mathbb{E}(X)$ and $\mathbb{E}(X) > 0$, then

$$\mathbb{P}[X \geq \mathbb{E}(X)] > 0, \quad (3.20)$$

and

$$\mathbb{P}[X > a] > 0 \text{ for } 0 \leq a < \mathbb{E}(X). \quad (3.21)$$

Further, for any real numbers μ , a and p satisfying $0 < a < \mu$ and $0 < p < 1$, there exists a nonnegative random variable Y such that

$$\mathbb{E}(Y) = \mu \text{ and } \mathbb{P}[Y \leq a] = p. \quad (3.22)$$

The above propositions show three things:

1. the Markov inequality is tight: if $0 < \mathbb{E}(X)/a < 1$, there is a nonnegative random variable X such that

$$\mathbb{P}[X \geq a] = \mathbb{E}(X)/a; \quad (3.23)$$

the inequality (3.2) cannot be improved without introducing further restrictions;

2. if $\mathbb{E}(X) > 0$, the probability $\mathbb{P}[X \geq a]$ can be arbitrarily large or small for any a in the interval $0 < a < \mathbb{E}(X)$;

3. the mean of X can be larger than any quantile of X , i.e. we can have $\mathbb{E}(X) > F_X^{-1}(q)$ for $0 < q < 1$.

The observations provide a way of illustrating the “ecological fallacy”.

Proposition 3.3 entails we can find, for any $0 < q < 1$, a random variable Y such that

$$\mathbb{E}(X) > F_X^{-1}(q). \quad (3.24)$$

Question – If X and Y are random variables, does the value of $\mathbb{E}(XY)$ impose restrictions on $\mathbb{P}[XY > a]$ or $\mathbb{P}[XY < a]$? For example, if $\mathbb{E}(XY) = \gamma > 0$, does this constrain $\mathbb{P}[XY > 0]$ or $\mathbb{P}[XY < 0]$?

3.3. Markov-type inequalities for bounded variables

When the random variable X has bounded support, both lower and upper bounds can be given for $\mathbb{P}[X \geq a]$ as follows.

3.4 Theorem TWO-SIDED MARKOV-TYPE INEQUALITIES FOR BOUNDED VARIABLES. Let X be a real random variable such that

$$\mathbb{P}[m \leq X \leq M] = 1 \quad (3.25)$$

where $-\infty \leq m \leq M \leq +\infty$. If X has finite expected value $\mathbb{E}(X)$, $a \in \mathbb{R}$ and $m < a < M$, then

$$\frac{\mathbb{E}(X) - a}{M - a} \leq \mathbb{P}[X > a] \leq \mathbb{P}[X \geq a] \leq \frac{\mathbb{E}(X) - m}{a - m}. \quad (3.26)$$

In Theorem 3.4, the positivity assumption is replaced by the more general support assumption (3.25), so X can take negative values. If $\mathbb{E}(X) = 0$ and $m \leq 0 \leq M$, we have:

$$\frac{-a}{M-a} \leq \mathbb{P}[X > a] \leq \mathbb{P}[X \geq a] \leq \frac{-m}{a-m}. \quad (3.27)$$

For example, for $m = -1$ and $M = 1$, we get:

$$\mathbb{P}[X > 0.5] \leq \mathbb{P}[X \geq 0.5] \leq \frac{1}{0.5+1} = \frac{1}{3}, \quad (3.28)$$

$$\mathbb{P}[X \geq -0.5] \geq \mathbb{P}[X > -0.5] \geq \frac{0.5}{1+0.5} = \frac{1}{3}. \quad (3.29)$$

The Markov inequality [Theorem 3.1] corresponds to the special case where $m = 0$ and $M = +\infty$. If $-\infty < m < M < +\infty$, the random variable X is degenerate at M , and $\mathbb{P}[X = M] = 1$.

For $m \geq 0$ and $m < \mathbb{E}(X) \leq a < M$, we also have

$$\frac{\mathbb{E}(X) - a}{M - a} \geq \frac{\mathbb{E}(X) - a}{M}, \quad \frac{\mathbb{E}(X) - m}{a - m} \leq \frac{\mathbb{E}(X)}{a} \quad (3.30)$$

so that

$$\frac{\mathbb{E}(X) - a}{M} \leq \mathbb{P}[X > a] \leq \mathbb{P}[X \geq a] \leq \frac{\mathbb{E}(X)}{a}. \quad (3.31)$$

The latter result follows from the *Basic inequality* of Loève (1977, Volume I, Section 9, p. 159) on taking $g(x) = x$. The bounds in (3.31) are however less tight than those in (3.26).

As in (3.3) - (3.7), we can see that (3.26) implies: for $m < a < M$,

$$\frac{a - \mathbb{E}(X)}{a - m} \leq \mathbb{P}[X < a] \leq \mathbb{P}[X \leq a] \leq \frac{M - \mathbb{E}(X)}{M - a}, \quad (3.32)$$

$$\mathbb{P}[X = a] \leq \min \left\{ \frac{\mathbb{E}(X) - m}{a - m}, \frac{M - \mathbb{E}(X)}{M - a} \right\}. \quad (3.33)$$

For example, for $a = 0$, we get:

$$\frac{\mathbb{E}(X)}{m} \leq \mathbb{P}[X < 0] \leq \mathbb{P}[X \leq 0] \leq \frac{M - \mathbb{E}(X)}{M}. \quad (3.34)$$

On taking $a = F_X^{-1}(q)$, we get for $0 < q < 1$:

$$\frac{F_X^{-1}(q) - \mathbb{E}(X)}{F_X^{-1}(q) - m} \leq \mathbb{P}[X < F_X^{-1}(q)] \leq \mathbb{P}[X \leq F_X^{-1}(q)] \leq \frac{M - \mathbb{E}(X)}{M - F_X^{-1}(q)} \quad (3.35)$$

hence

$$q \leq \mathbb{P}[X \leq F_X^{-1}(q)] \leq \frac{M - \mathbb{E}(X)}{M - F_X^{-1}(q)}, \quad (3.36)$$

$$\frac{F_X^{-1}(q) - \mathbb{E}(X)}{F_X^{-1}(q) - m} \leq \mathbb{P}[X < F_X^{-1}(q)] \leq q, \quad (3.37)$$

and

$$F_X^{-1}(q) - q[F_X^{-1}(q) - m] \leq \mathbb{E}(X) \leq M - q[M - F_X^{-1}(q)], \quad (3.38)$$

$$mq + (1 - q)F_X^{-1}(q) \leq \mathbb{E}(X) \leq qF_X^{-1}(q) + (1 - q)M. \quad (3.39)$$

Correspondingly,

$$\frac{\mathbb{E}(X) - (1 - q)M}{q} \leq F_X^{-1}(q) \leq \frac{\mathbb{E}(X) - mq}{(1 - q)}. \quad (3.40)$$

If $m = 0$, we have:

$$(1 - q)F_X^{-1}(q) \leq \mathbb{E}(X) \leq qF_X^{-1}(q) + (1 - q)M, \quad (3.41)$$

$$\frac{\mathbb{E}(X) - (1 - q)M}{q} \leq F_X^{-1}(q) \leq \frac{\mathbb{E}(X)}{(1 - q)}. \quad (3.42)$$

If $M = 0$, we have:

$$mq + (1 - q)F_X^{-1}(q) \leq \mathbb{E}(X) \leq qF_X^{-1}(q), \quad (3.43)$$

$$\frac{\mathbb{E}(X)}{q} \leq F_X^{-1}(q) \leq \frac{\mathbb{E}(X) - mq}{(1 - q)}. \quad (3.44)$$

Another advantage of (3.26) is the possibility of working in terms of deviation from the mean, for $X - \mathbb{E}(X)$ typically can take negative and positive values:

$$\frac{-a}{M - a} \leq \mathbb{P}[X - \mathbb{E}(X) > a] \leq \mathbb{P}[X - \mathbb{E}(X) \geq a] \leq \frac{-m}{a - m}, \quad (3.45)$$

$$\frac{a}{a - m} \leq \mathbb{P}[X - \mathbb{E}(X) < a] \leq \mathbb{P}[X - \mathbb{E}(X) \leq a] \leq \frac{M}{M - a}, \quad (3.46)$$

$$\mathbb{P}[X - \mathbb{E}(X) = a] \leq \min \left\{ \frac{-m}{a - m}, \frac{M}{M - a} \right\}. \quad (3.47)$$

3.4. Chebyshev-type inequalities

3.5 Theorem TWO-SIDED MONOTONIC MARKOV-TYPE INEQUALITIES. Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $g(X)$ is a real random variable, $\mathbb{E}(|g(X)|) < \infty$ and

$$\mathbb{P}[0 \leq g(X) \leq M] = 1 \quad (3.48)$$

where $M \in [0, \infty]$. If $g(x)$ is a non-decreasing function on \mathbb{R} , then, for all $a \in \mathbb{R}$ such that $g(a) > 0$,

$$\frac{\mathbb{E}[g(X)] - g(a)}{M} \leq \mathbb{P}[X \geq a] \leq \frac{\mathbb{E}[g(X)]}{g(a)}. \quad (3.49)$$

If $g(x)$ is a non-decreasing function on $[0, \infty)$ and $g(x) = g(-x)$ for any x , then, for all $a \geq 0$ such that $g(a) > 0$,

$$\frac{\mathbb{E}[g(X)] - g(a)}{M} \leq \mathbb{P}[|X| \geq a] \leq \frac{\mathbb{E}[g(X)]}{g(a)} \quad (3.50)$$

where $0/0 \equiv 1$.

3.6 Corollary CHEBYSHEV INEQUALITY. If $\mathbb{E}(|X|^r) < \infty$, for $r > 0$, and $\mathbb{P}[m \leq |X| \leq M] = 1$, where $0 \leq m \leq M < \infty$ and $M \leq \infty$, then, for all $a \in (m, M)$

$$\mathbb{P}[|X| \geq a] \leq \frac{\mathbb{E}(|X|^r)}{a^r}. \quad (3.51)$$

Note the above result does not require that X be positive. It is easy to see that (3.49) entails

$$\mathbb{P}[X \leq a] \geq \mathbb{P}[X < a] \geq 1 - \frac{\mathbb{E}[g(X)]}{g(a)} \quad (3.52)$$

while (3.50) entails

$$\mathbb{P}[|X| \leq a] \geq \mathbb{P}[|X| < a] \geq 1 - \frac{\mathbb{E}[g(X)]}{g(a)}. \quad (3.53)$$

On taking $g(x) = x^\alpha$, $\alpha > 0$, we see that the distribution function of X is dominated by the Pareto distribution

$$\begin{aligned} F_a(x) &= 1 - \frac{\mathbb{E}[X^\alpha]}{x^\alpha} && \text{for } x \geq \mathbb{E}[X^\alpha] \\ &= 0 && \text{for } x < \mathbb{E}[X^\alpha]. \end{aligned} \quad (3.54)$$

Take $M = \infty$, and suppose the function $g(x)$ is homogeneous of degree k , i.e.

$$g(\lambda x) = \lambda^k g(x) \text{ for all } x. \quad (3.55)$$

Then, (3.49) implies:

$$\mathbb{P}[X \geq a(\mathbb{E}[g(X)])] \leq \frac{\mathbb{E}[g(X)]}{\mathbb{E}[g(X)]^k g(a)} = \frac{\mathbb{E}[g(X)]^{1-k}}{g(a)}, \quad (3.56)$$

$$\mathbb{P}[X \geq a(\mathbb{E}[g(X)])^{1/k}] \leq \frac{\mathbb{E}[g(X)]}{\mathbb{E}[g(X)]g(a)} = \frac{1}{g(a)}. \quad (3.57)$$

hence for $g(x) = x^k$ and $a > 0$,

$$\mathbb{P}[X \geq a(\mathbb{E}[X^k])] \leq \frac{1}{a^k}, \quad (3.58)$$

$$\mathbb{P}[X \geq a(\mathbb{E}[X^k])^{1/k}] \leq \frac{1}{a^k}. \quad (3.59)$$

In particular,

$$\mathbb{P}[X \geq a\mathbb{E}[X]] \leq \frac{1}{a}, \quad (3.60)$$

$$\mathbb{P}[X \geq a(\mathbb{E}[X^2])^{1/2}] \leq \frac{1}{a^2}. \quad (3.61)$$

3.7 Theorem TWO-SIDED BOUNDED MONOTONIC MARKOV-TYPE INEQUALITIES. Let X be a real random variable, and $g : \mathbb{R} \rightarrow \mathbb{R}$ a function such that $g(X)$ is a real random variable which satisfies $\mathbb{E}(|g(X)|) < \infty$ and

$$\mathbb{P}[m \leq g(X) \leq M] = 1 \quad (3.62)$$

where $-\infty \leq m \leq M \leq +\infty$. If $g(x)$ is a non-decreasing function on \mathbb{R} , then, for all $a \in \mathbb{R}$ such that $m < g(a) < M$,

$$\frac{\mathbb{E}[g(X)] - g(a)}{M - g(a)} \leq \mathbb{P}[|X| > a] \leq \mathbb{P}[|X| \geq a] \leq \frac{\mathbb{E}[g(X)] - m}{g(a) - m}. \quad (3.63)$$

If $g(x)$ is a non-decreasing function on $[0, \infty)$ and $g(x) = g(-x)$ for any x , then, for all $a \geq 0$ such that $g(a) > 0$,

$$\frac{\mathbb{E}[g(X)] - g(a)}{M - g(a)} \leq \mathbb{P}[|X| > a] \leq \mathbb{P}[|X| \geq a] \leq \frac{\mathbb{E}[g(X)] - m}{g(a) - m} \quad (3.64)$$

where $0/0 \equiv 1$.

3.8 Corollary CHEBYSHEV INEQUALITIES. If $\mathbb{E}(|X|^r) < \infty$, for $r > 0$, and $\mathbb{P}[m \leq |X| \leq M] = 1$, where $0 \leq m \leq M \in, m < \infty$ and $M \leq \infty$, then, for all $a \in (m, M)$,

$$\frac{\mathbb{E}(|X|^r) - a^r}{M^r - a^r} \leq \mathbb{P}[|X| \geq a] \leq \frac{\mathbb{E}(|X|^r) - m^r}{a^r - m^r}. \quad (3.65)$$

3.5. Generalized Markov-type inequalities

3.9 Theorem GENERIC MARKOV-TYPE INEQUALITIES. Let Y be a real random variable and an event A such that

$$\mathbb{P}[A \cap \{m(A) \leq Y \leq M(A)\}] = \mathbb{P}(A), \quad (3.66)$$

$$\mathbb{P}[A^c \cap \{m(A^c) \leq Y \leq M(A^c)\}] = \mathbb{P}(A^c), \quad (3.67)$$

where $-\infty \leq m(A) \leq M(A) \leq +\infty$ and $-\infty \leq m(A^c) \leq M(A^c) \leq +\infty$. If Y has finite expected value $\mathbb{E}(Y)$, then

$$[m(A) - m(A^c)]\mathbb{P}(A) \leq \mathbb{E}(Y) - m(A^c), \quad (3.68)$$

$$[M(A^c) - M(A)]\mathbb{P}(A) \leq M(A^c) - \mathbb{E}(Y), \quad (3.69)$$

In particular,

$$\mathbb{P}(A) \leq \frac{\mathbb{E}(Y) - m(A^c)}{m(A) - m(A^c)}, \quad \text{if } m(A) > m(A^c), \quad (3.70)$$

$$\mathbb{P}(A) \geq \frac{\mathbb{E}(Y) - m(A^c)}{m(A) - m(A^c)}, \quad \text{if } m(A) < m(A^c), \quad (3.71)$$

$$\mathbb{P}(A) \leq \frac{M(A^c) - \mathbb{E}(Y)}{M(A^c) - M(A)}, \quad \text{if } M(A) < M(A^c), \quad (3.72)$$

$$\mathbb{P}(A) \geq \frac{M(A^c) - \mathbb{E}(Y)}{M(A^c) - M(A)}, \quad \text{if } M(A) > M(A^c). \quad (3.73)$$

Two-sided inequalities easily follow from Theorem 3.9. If $m(A) > m(A^c)$ and $M(A) > M(A^c)$, we have

$$\frac{M(A^c) - \mathbb{E}(Y)}{M(A^c) - M(A)} \leq \mathbb{P}(A) \leq \frac{\mathbb{E}(Y) - m(A^c)}{m(A) - m(A^c)}. \quad (3.74)$$

Similarly, if $m(A) < m(A^c)$ and $M(A) < M(A^c)$, we get:

$$\frac{\mathbb{E}(Y) - m(A^c)}{m(A) - m(A^c)} \leq \mathbb{P}(A) \leq \frac{M(A^c) - \mathbb{E}(Y)}{M(A^c) - M(A)}. \quad (3.75)$$

3.10 Theorem MARKOV-TYPE INEQUALITIES BASED ON TRUNCATED MOMENTS. Let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $g(X)$ is a real random variable, $\mathbb{E}(|g(X)|) < \infty$ and

$$0 \leq g(x) \leq M_U \text{ for } x \geq A_U, \quad (3.76)$$

$$0 \leq g(x) \leq M_L \text{ for } x \leq A_L, \quad (3.77)$$

where $0 \leq M_U \leq \infty$, $0 \leq M_L \leq \infty$, $0 \leq A_U \leq \infty$ and $0 \leq A_L \leq \infty$. Let also

$$C_U(g, a) = \int_{[a, \infty)} g(x) dF_X(x), \quad C_L(g, a) = \int_{(-\infty, a]} g(x) dF_X(x). \quad (3.78)$$

(a) If $g(x)$ is nondecreasing on $[A_U, \infty)$, then, for $a \geq A_U$,

$$\frac{C_U(g, a)}{M_U} \leq \mathbb{P}[X \geq a] \leq \frac{C_U(g, a)}{g(a)}. \quad (3.79)$$

(b) If $g(x)$ is nonincreasing on $(-\infty, A_L]$, then, for $a \leq A_L$,

$$\frac{C_L(g, a)}{M_L} \leq \mathbb{P}[X \leq a] \leq \frac{C_L(g, a)}{g(a)}. \quad (3.80)$$

(c) If $g(x)$ is nondecreasing on $[A_U, \infty)$ and nonincreasing on $(-\infty, A_L]$, then, for $a \geq \max\{|A_U|, |A_L|\}$,

$$\begin{aligned} \mathbb{P}[|X| \geq a] &\leq \frac{C_U(g, a)}{g(a)} + \frac{C_L(g, a)}{g(-a)} \\ &\leq \frac{C_U(g, a) + C_L(g, a)}{\min\{g(a), g(-a)\}}, \end{aligned} \quad (3.81)$$

$$\begin{aligned} \mathbb{P}[|X| \geq a] &\geq \frac{C_U(g, a)}{M_U} + \frac{C_L(g, a)}{M_L} \\ &\geq \frac{C_U(g, a) + C_L(g, a)}{\max\{M_U, M_L\}}. \end{aligned} \quad (3.82)$$

4. Moments as integrals of distribution and quantile functions

4.1 Proposition DISTRIBUTION DECOMPOSITION OF THE FIRST MOMENT. Let X be a random variable with distribution function $F_X(x)$. If $\mathbb{E}|X| < \infty$, then

$$\int_0^\infty xF_X(x) = \int_0^\infty \mathbb{P}[X \geq x]dx = \int_0^\infty [1 - F_X(x)]dx, \quad (4.1)$$

$$\int_{-\infty}^0 x dF_X(x) = - \int_{-\infty}^0 \mathbb{P}[X \leq x]dx = - \int_{-\infty}^0 F_X(x)dx, \quad (4.2)$$

$$\begin{aligned} \mathbb{E}(X) &= \int_0^\infty \mathbb{P}[X \geq x]dx - \int_{-\infty}^0 \mathbb{P}[X \leq x]dx, \\ &= \int_0^\infty [1 - F_X(x)]dx - \int_{-\infty}^0 F_X(x)dx \end{aligned} \quad (4.3)$$

$$= \int_0^\infty \{1 - F_X(x) - F_X(-x)\}dx. \quad (4.4)$$

4.2 Corollary DISTRIBUTION DECOMPOSITION OF THE FIRST ABSOLUTE MOMENT. Let X be a random variable with distribution function $F_X(x)$. If $\mathbb{E}|X| < \infty$, then

$$\begin{aligned} \mathbb{E}(|X|) &= \int_0^\infty [1 - F_X(x)]dx + \int_{-\infty}^0 F_X(x)dx \\ &= \int_0^\infty \mathbb{P}[X \geq x]dx + \int_{-\infty}^0 \mathbb{P}[X \leq x]dx \\ &= \int_{-\infty}^\infty \mathbb{P}[|X| \geq x]dx. \end{aligned} \quad (4.5)$$

4.3 Theorem DISTRIBUTION DECOMPOSITION OF THE EXPECTED VALUE OF $g(X)$. Let X be a real random variable, and $g : \mathbb{R} \rightarrow \mathbb{R}$ a function such that $g(X)$ is a real random variable which satisfies $\mathbb{E}(|g(X)|) < \infty$. If $g(x)$ is differentiable a.e. on \mathbb{R} , then, for any $a \in \mathbb{R}$,

$$\begin{aligned} \int_a^\infty g(x) dF_X(x) &= g(a)[1 - F_X(a)] + \int_a^\infty g'(x) [1 - F_X(x)] dx \\ &= g(a)\mathbb{P}[X > a] + \int_a^\infty g'(x)\mathbb{P}[X > x] dx \\ &= g(a)\mathbb{P}[X \geq a] + \int_a^\infty g'(x)\mathbb{P}[X \geq x] dx \\ &= \mathbb{P}[X \geq a]\{g(a) + \int_{-\infty}^\infty g'(x)\mathbb{P}[X \geq x | X \geq a] dx\} \end{aligned} \quad (4.6)$$

$$\begin{aligned}
\int_{-\infty}^a g(x) dF_X(x) &= g(a)F_X(a) - \int_{-\infty}^a g'(x)F_X(x) dx \\
&= g(a)\mathbb{P}[X \leq a] - \int_{-\infty}^a g'(x)\mathbb{P}[X \leq x] dx \\
&= \mathbb{P}[X \leq a] \{g(a) - \int_{-\infty}^a g'(x)\mathbb{P}[X \leq x | X \leq a] dx\},
\end{aligned} \tag{4.7}$$

$$\begin{aligned}
\mathbb{E}[g(X)] &= g(a) + \int_a^{\infty} g'(x)[1 - F_X(x)] dx - \int_{-\infty}^a g'(x)F_X(x) dx \\
&= g(a) + \int_a^{\infty} g'(x)\mathbb{P}[X > x] dx - \int_{-\infty}^a g'(x)\mathbb{P}[X \leq x] dx \\
&= g(a) + \mathbb{P}[X \geq a] \int_{-\infty}^a g'(x) \{ \mathbb{P}[X \geq x | X \geq a] dx \\
&\quad - \mathbb{P}[X \leq a] \int_{-\infty}^{\infty} g'(x)\mathbb{P}[X \leq x | X \leq a] dx \}.
\end{aligned} \tag{4.8}$$

It is of interest to spell out a number of special cases of (4.8):

$$\mathbb{E}[g(X)] = g(0) + \int_0^{\infty} g'(x)[1 - F_X(x)] dx - \int_{-\infty}^0 g'(x)F_X(x) dx \tag{4.9}$$

$$= g(0) + \int_0^{\infty} g'(x)[1 - F_X(x)] dx + \int_0^{\infty} g'(-x)F_X(-x) dx \tag{4.10}$$

$$= g(0) + \int_0^{\infty} \{g'(x)[1 - F_X(x)] + g'(-x)F_X(-x)\} dx \tag{4.11}$$

In particular, if $g(-x) = g(x)$, we have $g'(-x) = -g'(x)$, hence

$$\mathbb{E}[g(X)] = g(0) + \int_0^{\infty} g'(x)[1 - F_X(x)] dx - \int_0^{\infty} g'(x)F_X(-x) dx \tag{4.12}$$

$$= g(0) + \int_0^{\infty} g'(x)[1 - F_X(x) - F_X(-x)] dx \tag{4.13}$$

and, if X has a distribution symmetric about zero [$F_X(-x) = 1 - F_X(x)$], for all x],

$$\mathbb{E}[g(X)] = g(0). \tag{4.14}$$

$$\mathbb{E}[g(X)] - g(\mathbb{E}[X]) = \int_{\mathbb{E}[X]}^{\infty} g'(x)[1 - F_X(x)]dx - \int_{-\infty}^{\mathbb{E}[X]} g'(x)F_X(x)dx. \quad (4.15)$$

When $g(x)$ is convex, we see that

$$\mathbb{E}[g(X)] - g(\mathbb{E}[X]) = \int_{\mathbb{E}[X]}^{\infty} g'(x)[1 - F_X(x)]dx - \int_{-\infty}^{\mathbb{E}[X]} g'(x)F_X(x)dx \geq 0. \quad (4.16)$$

This yields a closed-form expression for the difference $\mathbb{E}[g(X)] - g(\mathbb{E}[X])$.

4.4 Corollary DISTRIBUTION DECOMPOSITION OF r -MOMENTS. Let X a random variable with distribution function $F_X(x)$ and $r > 0$. If $\mathbb{E}(|X|^r) < \infty$, then

$$\int_0^{\infty} x^r dF_X(x) = r \int_0^{\infty} x^{r-1} [1 - F_X(x)] dx, \quad (4.17)$$

$$\begin{aligned} \mathbb{E}(|X|^r) &= r \int_0^{\infty} x^{r-1} \mathbb{P}(|X| \geq x) dx \\ &= r \int_0^{\infty} x^{r-1} [1 - F_X(x) + F_X(-x)] dx. \end{aligned} \quad (4.18)$$

4.5 Proposition MOMENT-TAIL AREA INEQUALITIES. Let $g(x)$ be a nonnegative strictly increasing function on $[0, \infty)$ and let $g^{-1}(x)$ be the inverse function of g . Then,

$$\sum_{n=1}^{\infty} \mathbb{P}[|X| \geq g^{-1}(n)] \leq \mathbb{E}[g(X)] \leq \sum_{n=0}^{\infty} \mathbb{P}[|X| > g^{-1}(n)]. \quad (4.19)$$

In particular, for any $r > 0$,

$$\begin{aligned} \sum_{n=1}^{\infty} \mathbb{P}(|X| \geq n^{1/r}) &\leq \mathbb{E}(|X|^r) \leq \sum_{n=0}^{\infty} \mathbb{P}(|X| > n^{1/r}) \\ &\leq 1 + \sum_{n=1}^{\infty} \mathbb{P}(X > n^{1/r}). \end{aligned} \quad (4.20)$$

4.6 Corollary MEAN-TAIL AREA INEQUALITIES. If X is a positive random variable,

$$\sum_{n=1}^{\infty} \mathbb{P}(X \geq n) \leq \mathbb{E}(X) \leq 1 + \sum_{n=1}^{\infty} \mathbb{P}(X > n). \quad (4.21)$$

4.7 Proposition INTEGRAL OF THE PROBABILITY OF AN INTERVAL. Let X a random variable with distribution function $F_X(x)$ and $0 \leq a \leq b$. If $\mathbb{E}|X| < \infty$, then

$$\int \mathbb{P}[a < X \leq x + b] dx = \int [F_X(x + b) - F_X(x)] dx = b, \quad (4.22)$$

$$\int \mathbb{P}[x - a < X \leq x + b] dx = \int [F_X(x + b) - F_X(x - a)] dx = a + b. \quad (4.23)$$

4.8 Corollary INTEGRAL OF THE PROBABILITY OF A GENERAL INTERVAL. Let X a random variable with distribution function $F_X(x)$ and $a \leq b$. If $\mathbb{E}|X| < \infty$, then

$$\int \mathbb{P}[x + a < X \leq x + b] dx = \int [F_X(x + b) - F_X(x + a)] dx = b - a. \quad (4.24)$$

4.9 Proposition QUANTILE REPRESENTATION OF THE MEAN. Let X a random variable with distribution function $F_X(x)$ and quantile function

$$F_X^{-1}(q) = \inf\{x : F_X(x) \geq q\}, \quad 0 < q < 1. \quad (4.25)$$

If $\mathbb{E}|X| < \infty$, then

$$\mathbb{E}(X) = \int_0^1 F_X^{-1}(q) dq. \quad (4.26)$$

5. Integration by parts

In order to study conditions for the existence of moments, it will be useful to recall some results on integration by parts.

5.1. Standard results

5.1 Theorem EXISTENCE OF RIEMANN-STIELTJES INTEGRAL. Each one of the following conditions is sufficient for the existence of the Riemann-Stieltjes integral $\int_a^b g(x) df(x)$.

(a) $g(x)$ is continuous on $[a, b]$ and $f(x)$ is of bounded variation on $[a, b]$.

5.2 Theorem REPRESENTATION OF RIEMANN-STIELTJES INTEGRAL FOR CONTINUOUS-BV FUNCTIONS. If $g(x)$ is continuous on $[a, b]$ and $f(x)$ is of bounded variation on $[a, b]$, then

$$\int_a^b g(x) df(x) = \sum_{x \in J} g(x)[f(x+) - f(x-)] + \int_a^b g(x) df_c(x) \quad (5.1)$$

where J is the set of discontinuities of g on $[a, b]$,

$$f_c(x) = f(x) - f_s(x), \quad (5.2)$$

and

$$\begin{aligned} f_s(x) &= f(x) - f(x-) + \sum_{y \in J \cap [a, x]} [f(y+) - f(y-)] \quad \text{for } x \in (a, b] \\ &= 0 \quad \text{for } x = a \end{aligned} \tag{5.3}$$

$g_s(x)$ is called the *saltus function* of g on $[a, b]$.

5.3 Theorem INTEGRATION BY PARTS. *The Riemann-Stieltjes integral $\int_a^b g(x) df(x)$ exists if and only the Riemann-Stieltjes integral $\int_a^b f(x) dg(x)$ exists. Further, when $\int_a^b g(x) df(x)$ exists, we have:*

$$\int_a^b f(x) dg(x) + \int_a^b g(x) df(x) = f(b)g(b) - f(a)g(a). \tag{5.4}$$

5.2. Extensions

5.4 Lemma RIEMANN-STIELTJES INTEGRATION BY PARTS. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ two real-valued functions and $-\infty < a \leq b < +\infty$. If the (Riemann-Stieltjes) integral $\int_a^b g(x) df(x)$ exists, then the integrals $\int_a^b f(x) dg(x)$ and $\int_a^b [A - f(x)] dg(x)$ also exist and*

$$\begin{aligned} \int_a^b g(x) df(x) &= g(b)f(b) - g(a)f(a) - \int_a^b f(x) dg(x) \\ &= [A - f(a)]g(a) - g(b)[A - f(b)] + \int_a^b [A - f(x)] dg(x), \end{aligned} \tag{5.5}$$

for any real constant A , with

$$\int_a^b f(x) dg(x) = \int_a^b f(x) g'(x) dx \tag{5.6}$$

and

$$\int_a^b [A - f(x)] dg(x) = \int_a^b [A - f(x)] g'(x) dx \tag{5.7}$$

if g is continuous on $[a, b]$ as well as differentiable on (a, b) and the Riemann integral $\int_a^b f(x) g'(x) dx$ exists (where g' can take arbitrary real values at a and b).

5.5 Lemma CENTERED RIEMANN-STIELTJES INTEGRATION BY PARTS. *Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ two real-valued functions and $-\infty < a \leq c \leq b < +\infty$. If the integrals $\int_a^b g(x) df(x)$, $\int_a^c g(x) df(x)$ and $\int_c^b g(x) df(x)$ exist, then the integrals $\int_a^c f(x) dg(x)$ and $\int_c^b f(x) dg(x)$ also exist, and*

$$\begin{aligned} \int_a^b g(x) df(x) &= Ag(c) - \{g(b)[A - f(b)] + g(a)f(a)\} \\ &\quad + \int_c^b [A - f(x)] dg(x) - \int_a^c f(x) dg(x) \end{aligned} \tag{5.8}$$

for any real constant A , with

$$\int_a^c f(x) dg(x) = \int_a^c f(x)g'(x) dx \quad (5.9)$$

if g is continuous on $[a, c]$ as well as differentiable on (a, c) and the Riemann integral $\int_a^c f(x)g'(x)dx$ exists (where g' can take arbitrary real values at a and c), and

$$\int_c^b [A - f(x)] dg(x) = \int_c^b [A - f(x)]g'(x) dx \quad (5.10)$$

if g is continuous on $[c, b]$ as well as differentiable on (c, b) and the Riemann integral $\int_c^b [A - f(x)]g'(x)dx$ exists (where g' can take arbitrary real values at c and b).

5.6 Lemma IMPROPER RIEMANN-STIELTJES INTEGRATION BY PARTS. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ two real-valued functions, and $-\infty < a \leq c \leq b < +\infty$. Let us also denote

$$v_{-\infty}[gf] = \lim_{a \rightarrow -\infty} g(a)f(a), \quad v_{\infty}[gf] = \lim_{b \rightarrow \infty} g(b)f(b), \quad (5.11)$$

$$v_{-\infty}[g(1-f)] = \lim_{a \rightarrow -\infty} [1-f(a)]g(a), \quad v_{\infty}[g(1-f)] = \lim_{b \rightarrow \infty} g(b)[1-f(b)], \quad (5.12)$$

whenever the limit considered exists.

(a) If the (improper Riemann-Stieltjes) integral $\int_a^{+\infty} g(x)df(x)$ exists, then:

(i) $\int_a^{+\infty} f(x) dg(x)$ exists and

$$\int_a^{+\infty} g(x)df(x) = v_{\infty}[gf] - g(a)f(a) - \int_a^{+\infty} f(x) dg(x) \quad (5.13)$$

provided the limit $v_{\infty}[gf]$ exists;

(ii) $\int_a^{+\infty} [1-f(x)] dg(x)$ exists and

$$\int_a^{+\infty} g(x)df(x) = [1-f(a)]g(a) - v_{\infty}[g(1-f)] + \int_a^{+\infty} [1-f(x)] dg(x) \quad (5.14)$$

provided the limit $v_{\infty}[g(1-f)]$ exists.

(b) If the integral $\int_{-\infty}^b g(x)df(x)$ exists, then:

(i) $\int_{-\infty}^b f(x) dg(x)$ exists and

$$\int_{-\infty}^b g(x)df(x) = g(b)f(b) - v_{-\infty}[gf] - \int_{-\infty}^b f(x) dg(x) \quad (5.15)$$

provided the limit $v_{-\infty}[gf]$ exists;

(ii) $\int_{-\infty}^b [1 - f(x)] dg(x)$ exists and

$$\int_{-\infty}^b g(x) df(x) = v_{-\infty}[g(1-f)] - g(b)[1-f(b)] + \int_{-\infty}^b [1-f(x)] dg(x) \quad (5.16)$$

provided the limit $v_{-\infty}[g(1-f)]$ exists.

(c) If the integral $\int_{-\infty}^{+\infty} g(x) df(x)$ exists, then:

(i) $\int_{-\infty}^{+\infty} f(x) dg(x)$ exists and

$$\int_{-\infty}^{+\infty} g(x) df(x) = v_{\infty}[gf] - v_{-\infty}[gf] - \int_{-\infty}^{+\infty} f(x) dg(x), \quad (5.17)$$

provided the limits $v_{\infty}[gf]$ and $v_{-\infty}[gf]$ exist;

(ii) $\int_{-\infty}^{+\infty} [1-f(x)] dg(x)$ exists and

$$\int_{-\infty}^{+\infty} g(x) df(x) = v_{-\infty}[g(1-f)] - v_{\infty}[g(1-f)] + \int_{-\infty}^{+\infty} [1-f(x)] dg(x), \quad (5.18)$$

provided the limits $v_{-\infty}[g(1-f)]$ and $v_{\infty}[g(1-f)]$ exist.

Furthermore, $dg(x)$ may be replaced by $g'(x)dx$ whenever g is differentiable everywhere in interior of the interval of integration considered and the corresponding Riemann integral exists.

5.7 Lemma IMPROPER CENTERED RIEMANN-STIELTJES INTEGRATION BY PARTS. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ two real-valued functions and $-\infty < a \leq c \leq b < +\infty$.

(a) If the integrals $\int_a^{+\infty} g(x) df(x)$, $\int_a^c g(x) df(x)$ and $\int_c^{+\infty} g(x) df(x)$ exist, then the integrals $\int_a^c f(x) dg(x)$ and $\int_c^{+\infty} [1-f(x)] dg(x)$ also exist and

$$\begin{aligned} \int_a^{+\infty} g(x) df(x) &= g(c) - \{v_{\infty}[g(1-f)] + g(a)f(a)\} \\ &\quad + \int_c^{+\infty} [1-f(x)] dg(x) - \int_a^c f(x) dg(x), \end{aligned} \quad (5.19)$$

provided the limit $v_{\infty}[g(1-f)]$ exists.

(b) If the integrals $\int_{-\infty}^b g(x) df(x)$, $\int_{-\infty}^c g(x) df(x)$ and $\int_c^b g(x) df(x)$ exist, then the integrals $\int_{-\infty}^c f(x) dg(x)$ and $\int_c^b [1-f(x)] dg(x)$ also exist and

$$\begin{aligned} \int_{-\infty}^b g(x) df(x) &= g(c) - \{g(b)[1-f(b)] + v_{-\infty}[gf]\} \\ &\quad + \int_c^b [1-f(x)] dg(x) - \int_{-\infty}^c f(x) dg(x), \end{aligned} \quad (5.20)$$

provided the limit $v_{-\infty}[gf]$ exists.

(c) If the integrals $\int_{-\infty}^{+\infty} g(x)df(x)$, $\int_{-\infty}^c g(x)df(x)$ and $\int_c^{+\infty} g(x)df(x)$ exist, then the integrals $\int_{-\infty}^c f(x)dg(x)$ and $\int_c^{+\infty} [1-f(x)]dg(x)$ also exist and

$$\begin{aligned} \int_{-\infty}^{+\infty} g(x)df(x) &= g(c) - \{v_{\infty}[g(1-f)] + v_{-\infty}[gf]\} \\ &\quad + \int_c^{+\infty} [1-f(x)]dg(x) - \int_{-\infty}^c f(x)dg(x), \end{aligned} \quad (5.21)$$

provided the limits $v_{\infty}[g(1-f)]$ and $v_{-\infty}[gf]$ exist.

Furthermore, $dg(x)$ may be replaced by $g'(x)dx$ whenever g is differentiable everywhere in interior of the interval of integration considered and the corresponding Riemann integral exists.

5.8 Lemma BOUNDED MONOTONICITY CONDITION FOR TAIL CONVERGENCE OF AN INTEGRABLE FUNCTION. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ two real-valued functions, and let m, M be two real constants.

(a) If $f(x)$ is monotonic nondecreasing on the interval $(-\infty, m)$ with finite limit as $x \rightarrow -\infty$, and if g satisfies the inequality

$$|g(a)| \leq B_L(x), \text{ for } x \leq a < m \quad (5.22)$$

where $B_L(x)$ is a real-valued function such that $\int_{-\infty}^m B_L(x)df(x)$ exists, then

$$0 \leq |g(a)|[f(a) - f(-\infty)] \leq \int_{-\infty}^a B_L(x)df(x), \text{ for } a < m, \quad (5.23)$$

where $f(-\infty) = \lim_{x \rightarrow -\infty} f(x) > -\infty$, and

$$\lim_{a \rightarrow -\infty} g(a)[f(a) - f(-\infty)] = 0. \quad (5.24)$$

(b) If $f(x)$ is monotonic nondecreasing on the interval $[M, \infty)$ with finite limit as $x \rightarrow \infty$, and if g satisfies the inequality

$$|g(b)| \leq |g(x)| + B_U(x), \text{ for } x \geq b > M \quad (5.25)$$

where $B_U(x)$ is a real-valued function such that $\int_M^{\infty} B_U(x)df(x)$ exists, then

$$0 \leq |g(b)|[f(\infty) - f(b)] \leq \int_b^{\infty} B_U(x)df(x), \text{ for } b > M, \quad (5.26)$$

where $f(\infty) = \lim_{x \rightarrow \infty} f(x) < \infty$, and

$$\lim_{b \rightarrow \infty} g(b) [f(\infty) - f(b)] = 0. \quad (5.27)$$

It is easy to see that (5.22) holds whenever $\int_{-\infty}^m |g(x)| df(x)$ exists and one of the following conditions holds: for some real constant B ,

$$|g(a)| \leq |g(x)| + B, \text{ for } x \leq a < m; \quad (5.28)$$

$$|g(x)| \text{ is nondecreasing on the interval } (M, \infty); \quad (5.29)$$

$$g(x) \text{ is bounded on the interval } (M, \infty). \quad (5.30)$$

Further, in case (5.28), we have:

$$0 \leq |g(a)| [f(a) - f(-\infty)] \leq \int_{-\infty}^a |g(x)| df(x) + B [f(a) - f(-\infty)]. \quad (5.31)$$

Similarly, (5.25) holds whenever one of the following conditions holds: for some real constant B ,

$$|g(a)| \leq |g(x)| + B, \text{ for } x > b > M; \quad (5.32)$$

$$|g(x)| \text{ is nonincreasing on the interval } (-\infty, m); \quad (5.33)$$

$$g(x) \text{ is bounded on the interval } (-\infty, m). \quad (5.34)$$

Further, in case (5.32), we have:

$$0 \leq |g(b)| [f(\infty) - f(b)] \leq \int_b^{\infty} |g(x)| df(x) + B [f(\infty) - f(b)] \quad (5.35)$$

6. Tail decay rates and the existence of moments

It follows from Markov-type inequalities in Theorem 3.7 that tail probabilities decay to zero at least as fast as $g(x)$ or x^k increases. More precisely, on taking $m = 0$, we see from Theorem 3.7, that

$$g(a) \mathbb{P}[|X| \geq a] \leq \mathbb{E}[g(X)] \text{ for all } a > 0 \quad (6.1)$$

hence

$$\limsup_{a \rightarrow \infty} \{g(a) \mathbb{P}[|X| \geq a]\} = \limsup_{a \rightarrow \infty} \left\{ \frac{\mathbb{P}[|X| \geq a]}{1/g(a)} \right\} \leq \mathbb{E}[g(X)]; \quad (6.2)$$

the rate of convergence of $\mathbb{P}[|X| \geq a]$ to zero must be at least as fast as the one of $1/g(a)$. In particular, if $\mathbb{E}(|X|^r) < \infty$ and $m = 0$,

$$a^r \mathbb{P}[|X| \geq a] \leq \mathbb{E}(|X|^r) \text{ for all } a > 0 \quad (6.3)$$

hence

$$\limsup_{a \rightarrow \infty} \{a^r \mathbb{P}[|X| \geq a]\} = \limsup_{a \rightarrow \infty} \left\{ \frac{\mathbb{P}[|X| \geq a]}{1/a^r} \right\} \leq \mathbb{E}(|X|^r); \quad (6.4)$$

the rate of convergence of $\mathbb{P}[|X| \geq a]$ to zero must be at least as fast as the one of $1/a^r$.

However, it is possible to make stronger statements on these rates of convergence by considering further generalizations of the Markov inequality.

6.1 Proposition TAIL DECAY RATES BASED ON TRUNCATED MOMENTS. *Under the assumptions of Theorem 3.10, the following limits hold:*

(a) *if $g(x)$ is nondecreasing on $[A_U, \infty)$, and if $g(a) > 0$ for $a \geq A_U$, then*

$$\limsup_{a \rightarrow \infty} \{g(a) \mathbb{P}[X \geq a]\} = \limsup_{a \rightarrow \infty} \left\{ \frac{\mathbb{P}[X \geq a]}{1/g(a)} \right\} = 0; \quad (6.5)$$

(b) *if $g(x)$ is nonincreasing on $(-\infty, A_L]$, and if $g(a) > 0$ for $a \leq A_L$, then,*

$$\limsup_{a \rightarrow -\infty} \{g(a) \mathbb{P}[X \leq a]\} = \limsup_{a \rightarrow -\infty} \left\{ \frac{\mathbb{P}[X \leq a]}{1/g(a)} \right\} = 0; \quad (6.6)$$

(c) *if $g(x)$ is nondecreasing on $[A_U, \infty)$ and nonincreasing on $(-\infty, A_L]$, and if $g(a) > 0$ for $|a| \geq \max\{|A_U|, |A_L|\}$, then*

$$\limsup_{a \rightarrow \infty} \{g(a) \mathbb{P}[|X| \geq a]\} = \limsup_{a \rightarrow \infty} \left\{ \frac{\mathbb{P}[|X| \geq a]}{1/g(a)} \right\} = 0. \quad (6.7)$$

Note that the limits (6.5) - (6.7), can be rewritten using the “little oh” notation $o(\cdot)$ [see Serfling (1980, Section 1.1.2)]:

$$\mathbb{P}[X \geq a] = o(g(a)^{-1}), \quad a \rightarrow \infty, \quad (6.8)$$

$$\mathbb{P}[X \leq a] = o(g(a)^{-1}), \quad a \rightarrow -\infty, \quad (6.9)$$

$$\mathbb{P}[|X| \geq a] = o(g(a)^{-1}), \quad a \rightarrow \infty. \quad (6.10)$$

In particular, if $g(x) = |x|^r$, we have:

$$\mathbb{P}[X \geq a] = o(|a|^{-r}), \quad a \rightarrow \infty, \quad (6.11)$$

$$\mathbb{P}[X \leq a] = o(|a|^{-r}), \quad a \rightarrow -\infty, \quad (6.12)$$

$$\mathbb{P}[|X| \geq a] = o(|a|^{-r}), \quad a \rightarrow \infty. \quad (6.13)$$

6.2 Corollary MOMENT EXISTENCE AND TAIL AREA DECAY. Let $r > 0$. If $\mathbb{E}(|X|^r) < \infty$, then

$$\begin{aligned} \lim_{x \rightarrow \infty} \{x^r \mathbb{P}[X \geq x]\} &= \lim_{x \rightarrow \infty} \{|x|^r \mathbb{P}[X \leq x]\} \\ &= \lim_{x \rightarrow \infty} \{x^r \mathbb{P}[|X| \geq x]\} = 0. \end{aligned} \quad (6.14)$$

In particular, if $\mathbb{E}(|X|) < \infty$, then

$$\begin{aligned} \lim_{x \rightarrow \infty} \{x \mathbb{P}[X \geq x]\} &= \lim_{x \rightarrow \infty} \{|x| \mathbb{P}[X \leq x]\} \\ &= \lim_{x \rightarrow \infty} \{x \mathbb{P}[|X| \geq x]\} = 0. \end{aligned} \quad (6.15)$$

The following proposition provides a general sufficient condition for the existence of a finite mean in terms of tail areas. It is a direct consequence of Proposition 4.1.

6.3 Proposition NECESSARY AND SUFFICIENT CONDITION FOR THE EXISTENCE OF THE MEAN. Let X be a random variable with distribution function $F_X(x)$. Then

$$\begin{aligned} \mathbb{E}(|X|) < \infty &\Leftrightarrow \mathbb{P}(|X| \geq x) \text{ is integrable on } (0, +\infty) \\ &\Leftrightarrow [1 - F_X(x) - F_X(-x)] \text{ is integrable on } (0, +\infty) \\ &\Leftrightarrow \int_0^\infty [1 - F_X(x)] dx < \infty \text{ and } \int_{-\infty}^0 F_X(x) dx < \infty. \end{aligned} \quad (6.16)$$

6.4 Proposition NECESSARY AND SUFFICIENT CONDITION FOR THE EXISTENCE OF r -MOMENTS. Let X a random variable with distribution function $F_X(x)$ and $r > 0$. If $\mathbb{E}(|X|^r) < \infty$, then

$$\begin{aligned} \mathbb{E}(|X|^r) < \infty &\Leftrightarrow x^{r-1} \mathbb{P}(|X| \geq x) \text{ is integrable on } (0, +\infty) \\ &\Leftrightarrow |x|^{r-1} [1 - F_X(x) + F_X(-x)] \text{ is integrable on } (0, +\infty) \\ &\Leftrightarrow \int_0^\infty x^{r-1} [1 - F_X(x)] dx < \infty \text{ and } \int_{-\infty}^0 |x|^{r-1} F_X(x) dx < \infty. \end{aligned} \quad (6.17)$$

7. Moments of sums of random variables

In this section, we consider a sequence X_1, \dots, X_n of random variables, and study the moments of the corresponding sum and average:

$$S_n = \sum_{i=1}^n X_i, \quad \bar{X}_n = S_n/n. \quad (7.1)$$

7.1 Proposition BOUNDS ON THE ABSOLUTE MOMENTS OF A SUM OF RANDOM VARIABLES.

$$\begin{aligned} \mathbb{E}(|S_n|^r) &\leq \sum_{i=1}^n \mathbb{E}(|X_i|^r), & \text{if } 0 < r \leq 1, \\ &\leq n^{r-1} \sum_{i=1}^n \mathbb{E}(|X_i|^r), & \text{if } r > 1, \end{aligned} \tag{7.2}$$

and

$$\begin{aligned} \mathbb{E}(|\bar{X}_n|^r) &\leq \left(\frac{1}{n}\right)^r \sum_{i=1}^n \mathbb{E}(|X_i|^r), & \text{if } 0 < r \leq 1, \\ &\leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}(|X_i|^r), & \text{if } r > 1. \end{aligned} \tag{7.3}$$

7.2 Proposition MINKOWSKI INEQUALITY FOR n VARIABLES. If $r \geq 1$, then

$$[\mathbb{E}(|S_n|^r)]^{1/r} \leq \sum_{i=1}^n [\mathbb{E}(|X_i|^r)]^{1/r} \tag{7.4}$$

and

$$\begin{aligned} [\mathbb{E}(|\bar{X}_n|^r)]^{1/r} &\leq \frac{1}{n} \sum_{i=1}^n [\mathbb{E}(|X_i|^r)]^{1/r} \\ &\leq \left\{ \frac{1}{n} \sum_{i=1}^n \mathbb{E}(|X_i|^r) \right\}^{1/r}. \end{aligned} \tag{7.5}$$

7.3 Proposition BOUNDS ON THE ABSOLUTE MOMENTS OF A SUM OF RANDOM VARIABLES UNDER CONDITIONAL SYMMETRY. If the distribution of X_{k+1} given S_k is symmetric about zero for $k = 1, \dots, n-1$, and $\mathbb{E}(|X_i|^r) < \infty$, $i = 1, \dots, n$, then

$$\mathbb{E}(|S_n|^r) \leq \sum_{i=1}^n \mathbb{E}(|X_i|^r) \quad \text{for } 1 \leq r \leq 2, \tag{7.6}$$

and

$$\mathbb{E}(|\bar{X}_n|^r) \leq \left(\frac{1}{n}\right)^r \sum_{i=1}^n \mathbb{E}(|X_i|^r) \quad \text{for } 1 \leq r \leq 2, \tag{7.7}$$

with equality holding when $r = 2$.

7.4 Proposition BOUNDS ON THE ABSOLUTE MOMENTS OF A SUM OF RANDOM VARIABLES UNDER MARTINGALE CONDITION. *If*

$$\mathbb{E}(X_{k+1} | S_k) = 0 \quad \text{a.s.,} \quad k = 1, \dots, n-1, \quad (7.8)$$

and $\mathbb{E}(|X_i|^r) < \infty, i = 1, \dots, n$, then

$$\mathbb{E}(|S_n|^r) \leq 2 \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad \text{for } 1 \leq r \leq 2, \quad (7.9)$$

and

$$\mathbb{E}(|\bar{X}_n|^r) \leq 2 \left(\frac{1}{n}\right)^r \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad \text{for } 1 \leq r \leq 2. \quad (7.10)$$

Furthermore, for $r = 2$,

$$\mathbb{E}(S_n^2) = \sum_{i=1}^n \mathbb{E}(X_i^2). \quad (7.11)$$

7.5 Proposition BOUNDS ON THE ABSOLUTE MOMENTS OF A SUM OF RANDOM VARIABLES UNDER TWO-SIDED MARTINGALE CONDITION. *Let*

$$S_{m(k)} = \sum_{i=1, i \neq k}^{m+1} X_i, \quad 1 \leq k \leq m+1 \leq n. \quad (7.12)$$

If

$$\mathbb{E}(X_k | S_{m(k)}) = 0 \quad \text{a.s.,} \quad \text{for } 1 \leq k \leq m+1 \leq n, \quad (7.13)$$

and $\mathbb{E}(|X_i|^r) < \infty, i = 1, \dots, n$, then

$$\mathbb{E}(|S_n|^r) \leq \left(2 - \frac{1}{n}\right)^r \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad \text{for } 1 \leq r \leq 2, \quad (7.14)$$

and

$$\mathbb{E}(|\bar{X}_n|^r) \leq \left(\frac{1}{n}\right)^r \left(2 - \frac{1}{n}\right)^r \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad \text{for } 1 \leq r \leq 2. \quad (7.15)$$

7.6 Proposition BOUNDS ON THE ABSOLUTE MOMENTS OF A SUM OF INDEPENDENT RANDOM VARIABLES. *Let the random variables X_1, \dots, X_n be independent with $\mathbb{E}(X_i) = 0$ and $\mathbb{E}(|X_i|^r) < \infty, i = 1, \dots, n$, and let*

$$D(r) = [13.52 / (2.6\pi)^r] \Gamma(r) \sin(r\pi/2). \quad (7.16)$$

If $D(r) < 1$ and $1 \leq r \leq 2$, then

$$\mathbb{E}(|S_n|^r) \leq [1 - D(r)]^{-1} \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad (7.17)$$

and

$$\mathbb{E}(|\bar{X}_n|^r) \leq \left(\frac{1}{n}\right)^r [1 - D(r)]^{-1} \sum_{i=1}^n \mathbb{E}(|X_i|^r), \quad \text{for } 1 \leq r \leq 2. \quad (7.18)$$

8. Proofs and references

1.1 to 3.8. See Loève (1977, Volume I, Sections 9.1 and 9.3, pp. 151-162). For Jensen inequality, see also Chow and Teicher (1988, Section 4.3, pp. 103-106). Hannan (1985), Lehmann and Shaffer (1988), Piegorsch and Casella (1988) and Khuri and Casella (2002) discussed conditions for the existence of the moments of $1/X$.

2.9. See von Bahr and Esseen (1965, Lemma 3).

3.10

PROOF OF THEOREM 3.10 (a) For $x \geq a \geq A_U$, we have $g(x) \geq g(a)$ and $g(x) \leq M_U$, hence

$$C_U(g, a) = \int_{[a, \infty)} g(x) dF_X(x) \geq g(a) \int_{[a, \infty)} dF_X(x) = g(a) \mathbb{P}[X \geq a]$$

and

$$\int_{[a, \infty)} g(x) dF_X(x) \leq M_U \mathbb{P}[X \geq a],$$

from which we get the inequality

$$\frac{C_U(g, a)}{M_U} \leq \mathbb{P}[X \geq a] \leq \frac{C_U(g, a)}{g(a)}.$$

(b) For $x \leq a \leq A_L$, we have $g(x) \geq g(a)$ and $g(x) \leq M_L$, hence

$$C_L(g, a) = \int_{[-\infty, a]} g(x) dF_X(x) \geq g(a) \int_{[-\infty, a]} dF_X(x) = g(a) \mathbb{P}[X \leq a]$$

and

$$\int_{[-\infty, a]} g(x) dF_X(x) \leq M_L \mathbb{P}[X \leq a]$$

from which we get the inequality

$$\frac{C_L(g, a)}{M_L} \leq \mathbb{P}[X \leq a] \leq \frac{C_L(g, a)}{g(a)}.$$

(c) For $a \geq \max(|A_U|, |A_L|)$, we have $a \geq A_U$ and $-a \leq A_L$, hence

$$\begin{aligned} \mathbb{P}\mathbb{P}[|X| \geq a] &= \mathbb{P}[X \geq a] + \mathbb{P}[X \leq -a] \\ &\leq \frac{C_U(g, a)}{g(a)} + \frac{C_L(g, a)}{g(a)} \end{aligned}$$

$$\leq \frac{C_U(g, a) + C_L(g, a)}{\min\{g(a), g(-a)\}}$$

and

$$\begin{aligned} \mathbb{P}[|X| \geq a] &\geq \frac{C_U(g, a)}{M_U} + \frac{C_L(g, a)}{M_L} \\ &\geq \frac{C_U(g, a) + C_L(g, a)}{\max(M_U, M_L)}. \end{aligned}$$

□

PROOF OF PROPOSITION 4.1 We have:

$$\mathbb{E}(X) = \int_0^\infty x dF_X(x) + \int_{-\infty}^0 x dF_X(x). \quad (8.1)$$

For any statement p , let us set

$$I(p) = \begin{cases} 1 & \text{if } p \text{ is true} \\ 0 & \text{if } p \text{ is false} \end{cases}. \quad (8.2)$$

Using this notation, we can write:

$$\begin{aligned} x &= \int I(0 < y < x) dy && \text{for } x \geq 0 \\ &= - \int I(x \leq y \leq 0) dy && \text{for } x < 0 \end{aligned}. \quad (8.3)$$

Then,

$$\begin{aligned} \int_0^\infty x F_X(x) &= \int I(x > 0) x dF_X(x) \\ &= \int I(x > 0) \left[\int I(0 < y < x) dy \right] dF_X(x) \\ &= \int \int I(x > 0) I(0 < y < x) dF_X(x) dy \\ &= \int \int I(y < x) I(y > 0) dF_X(x) dy \\ &= \int \left[\int I(y < x) dF_X(x) \right] I(y > 0) dy \\ &= \int \left[\int_{x>y} dF_X(x) \right] I(y > 0) dy \end{aligned}$$

$$\begin{aligned}
&= \int [1 - F_X(y)]I(y > 0)dy = \int_0^\infty [1 - F_X(y)]dy \\
&= \int_0^\infty [1 - F_X(x)]dx = \int_0^\infty \mathbb{P}[X \geq x]dx.
\end{aligned} \tag{8.4}$$

Similarly,

$$\begin{aligned}
\int_{-\infty}^0 xF_X(x)dx &= \int I(x \leq 0)x dF_X(x) \\
&= - \int I(x \leq 0) \left[\int I(x \leq y \leq 0)dy \right] dF_X(x) \\
&= - \int \int I(x \leq y \leq 0)I(x \leq 0) dF_X(x) dy \\
&= - \int \int I(x \leq y)I(y \leq 0) dF_X(x) dy \\
&= - \int \left[\int I(x \leq y) dF_X(x) \right] I(y \leq 0) dy \\
&= - \int F_X(y)I(y \leq 0) dy = - \int_{-\infty}^0 F_X(y) dy \\
&= - \int_{-\infty}^0 F_X(x) dx = - \int_{-\infty}^0 \mathbb{P}[X \leq x] dx.
\end{aligned} \tag{8.5}$$

Therefore,

$$\begin{aligned}
\mathbb{E}(X) &= \int_0^\infty [1 - F_X(x)]dx - \int_{-\infty}^0 F_X(x)dx \\
&= \int_0^\infty \mathbb{P}[X \geq x]dx - \int_{-\infty}^0 \mathbb{P}[X \leq x]dx.
\end{aligned} \tag{8.6}$$

Finally, (4.5) follows on observing that

$$\int_{-\infty}^0 |x| F_X(x) dx = - \int_{-\infty}^0 x F_X(x) dx = \int_{-\infty}^0 F_X(x) dx \int_{-\infty}^0 \mathbb{P}[X \leq x] dx. \tag{8.7}$$

□

4.2 to 6.2. See Feller (1966, Section V.6, Lemma 1), Chung (1974, Section 3.2, Exercises 17-18), Serfling (1980, Section 1.14, pp. 46-47) and Chow and Teicher (1988,

Section 4.3, pp. 103-106). For other inequalities involving absolute moments, the reader may consult Beesack (1984). Further discussion of Proposition 4.1 is available in Hong (2012, 2015) and Lo (2019).

PROOF OF THEOREM 4.3 For any $a \in \mathbb{R}$, we have:

$$\mathbb{E}[g(X)] = \int_a^\infty g(x) dF_X(x) + \int_{-\infty}^a g(x) dF_X(x). \quad (8.8)$$

For any statement p , let us set

$$I(p) = \begin{cases} 1 & \text{if } p \text{ is true} \\ 0 & \text{if } p \text{ is false.} \end{cases} \quad (8.9)$$

By the fundamental theorem of calculus, we can write: for any $a \in \mathbb{R}$,

$$g(x) = \begin{cases} g(a) + \int_a^x g'(y) dy & \text{if } x \geq a \\ g(a) - \int_a^x g'(y) dy & \text{if } x < a \end{cases}. \quad (8.10)$$

Then,

$$\begin{aligned} \int_a^\infty g(x) dF_X(x) &= \int I(a < x) g(x) dF_X(x) \\ &= \int I(a < x) [g(a) + \int_a^x g'(y) dy] dF_X(x) \\ &= \int I(a < x) g(a) dF_X(x) + \int I(a < x) \left[\int_a^x g'(y) dy \right] dF_X(x) \\ &= g(a) \mathbb{P}[X > a] + \int I(a < x) \left[\int I(a < y \leq x) g'(y) dy \right] dF_X(x) \\ &= g(a) \mathbb{P}[X > a] + \iint I(a < x) I(a < y \leq x) g'(y) dy dF_X(x) \\ &= g(a) \mathbb{P}[X > a] + \iint I(y \leq x) I(a < y) g'(y) dy dF_X(x) \\ &= g(a) \mathbb{P}[X > a] + \iint I(a < y) I(y \leq x) dF_X(x) dy \\ &= g(a) \mathbb{P}[X > a] + \int I(a < y) g'(y) \left[\int I(y \leq x) dF_X(x) \right] dy \\ &= g(a) \mathbb{P}[X > a] + \int I(a < y) g'(y) \mathbb{P}[X \geq y] dy \\ &= g(a) \mathbb{P}[X > a] + \int I(a < y) g'(y) \mathbb{P}[X > y] dy \end{aligned}$$

$$\begin{aligned}
&= g(a)\mathbb{P}[X > a] + \int_a^\infty g'(y)\mathbb{P}[X > y]dy \\
&= g(a)\mathbb{P}[X > a] + \int_a^\infty g'(y)[1 - F_X(y)]dy.
\end{aligned} \tag{8.11}$$

Similarly,

$$\begin{aligned}
\int_{-\infty}^u g(x)dF_X(x) &= \int I(x \leq a)g(x)dF_X(x) \\
&= \int I(x \leq a)[g(a) - \int_a^x g'(y)dy]dF_X(x) \\
&= g(a) \int I(x \leq a)dF_X(x) - \int I(x \leq a) \left[\int_a^x g'(y)dy \right] dF_X(x) \\
&= g(a)\mathbb{P}[X \leq a] - \int I(x \leq a) \left[\int I(x < y \leq a)g'(y)dy \right] dF_X(x) \\
&= g(a)\mathbb{P}[X \leq a] - \int \int I(x \leq a)I(x < y \leq a)g'(y)dydF_X(x) \\
&= g(a)\mathbb{P}[X \leq a] - \int \int I(x < y)I(y \leq a)g'(y)dydF_X(x) \\
&= g(a)\mathbb{P}[X \leq a] - \int \int I(y \leq a)g'(y)I(x < y)dF_X(x)dy \\
&= g(a)\mathbb{P}[X \leq a] - \int I(y \leq a)g'(y) \left[\int I(x < y)dF_X(x) \right] dy \\
&= g(a)\mathbb{P}[X \leq a] - \int I(y \leq a)g'(y)\mathbb{P}[X < y]dy \\
&= g(a)\mathbb{P}[X \leq a] - \int I(y \leq a)g'(y)\mathbb{P}[X \leq y]dy \\
&= g(a)\mathbb{P}[X \leq a] - \int_{-\infty}^u g'(y)\mathbb{P}[X \leq y]dy \\
&= g(a)F_X(a) - \int_{-\infty}^u g'(y)F_X(y)dy.
\end{aligned} \tag{8.12}$$

Therefore,

$$\mathbb{E}[g(X)] = \int_a^\infty g(x)dF_X(x) + \int_{-\infty}^u g(x)dF_X(x)$$

$$\begin{aligned}
&= g(a) + \int_a^\infty g'(y) [1 - F_X(y)] dy - \int_{-\infty}^a g'(y) F_X(y) dy \\
&= g(a) + \int_a^\infty g'(y) \mathbb{P}[X > y] dy - \int_{-\infty}^a g'(y) \mathbb{P}[X \leq y] dy \\
&= g(a) + \int_a^\infty g'(y) \mathbb{P}[X \geq y] dy - \int_{-\infty}^a g'(y) \mathbb{P}[X \leq y] dy
\end{aligned} \tag{8.13}$$

□

4.7 The identity (4.22) is stated by Chung (1974, Section 3.2, Exercise 16). We give below a simple proof along with a slight extension. PROOF OF PROPOSITION 4.7 We can establish (4.22) as follows, through the use of the indicator function $I(p)$ defined in (8.2):

$$\begin{aligned}
\int [F_X(x+b) - F_X(x)] dx &= \int \mathbb{P}[a < X \leq x+b] dx \\
&= \int [\int I(x < y \leq x+b) dF_X(y)] dx \\
&= \int [\int I(x < y \leq x+b) dx] dF_X(y) \\
&= \int b dF_X(y) = b.
\end{aligned} \tag{8.14}$$

(4.23) then follows by repeated application of the above identity:

$$\begin{aligned}
\int [F_X(x+b) - F_X(x-a)] dx &= \int \mathbb{P}[x-a < X \leq x+b] dx \\
&= \int \mathbb{P}[x-a < X \leq x] dx + \int \mathbb{P}[x < X \leq x+b] dx \\
&= \int \mathbb{P}[x < X \leq x+a] dx + \int \mathbb{P}[x < X \leq x+b] dx \\
&= a+b.
\end{aligned} \tag{8.15}$$

□

4.5. See Chow and Teicher (1988, Section 4.1, Corollary 3, p. 90).

4.6. The inequality (4.21) is given by Chung (1974, Theorem 3.2.1) and Serfling (1980, Section 1.3, p. 12). Theorem 5.1 (a) See Riesz and Sz.-Nagy (1955/1990, Section 54) and Devinatz (1968, Theorem 5.5.2, page 219).

Theorem 5.2. See Devinatz (1968, Theorem 5.5.7, page 225).

Theorem 5.3 See Riesz and Sz.-Nagy (1955/1990, Section 54) and Haaser and Sullivan (1991, Theorem 2.8, page 254).

Theorem 5.4 See Riesz and Sz.-Nagy (1955/1990, Section 54) for $A = 0$.

PROOF OF LEMMA 5.4 The first identity in (5.5) part is given by Devinatz (1968, Theorem 5.4.8, page 213) and Protter and Morrey (1991, Theorem 12.12, page 320), while the second follows from the latter on observing that

$$\begin{aligned} -\int_a^b f(x) dg(x) &= \int_a^b [A - f(x) - A] dg(x) = \int_a^b [A - f(x)] dg(x) - A \int_a^b dg(x) \\ &= \int_a^b [A - f(x)] dg(x) - A [g(b) - g(a)] \end{aligned} \quad (8.16)$$

and rearranging the terms of the sum. Equation (8.16) also entails the existence of the integral $\int_a^b [1 - f(x)] dg(x)$. The identities (5.6)-(5.7) follow on observing that we can write $dg(x) = g'(x)dx$ when g is differentiable [see Devinatz (1968, Theorem 5.4.7, page 213)]. \square

5.5

PROOF OF LEMMA 5.5 Using Lemma 5.5, we get:

$$\begin{aligned} \int_a^b g(x) df(x) &= \int_a^c g(x) df(x) + \int_c^b g(x) df(x) \\ &= g(c)f(c) - g(a)f(a) - \int_a^c f(x) dg(x) \\ &\quad + [A - f(c)]g(c) - g(b)[A - f(b)] + \int_c^b [A - f(x)] dg(x) \\ &= Ag(c) - \{g(b)[A - f(b)] + g(a)f(a)\} \\ &\quad + \int_c^b [A - f(x)] dg(x) - \int_a^c f(x) dg(x). \end{aligned} \quad (8.17)$$

\square

5.8

PROOF OF LEMMA 5.8 (a) The existence of the limit $\lim_{x \rightarrow -\infty} f(x)$ entails that the integral $\int_{-\infty}^a df(x) = f(a) - f(-\infty)$ also exists. Since $f(x)$ is monotonic nondecreasing on the interval $(-\infty, m)$ and $\int_{-\infty}^m B_L(x) df(x)$ exists, we get from (5.22): for $a < m$,

$$0 \leq \int_{-\infty}^a |g(a)| df(x) \leq \int_{-\infty}^a B_L(x) df(x) \quad (8.18)$$

hence

$$0 \leq |g(a)| [f(a) - f(-\infty)] \leq \int_{-\infty}^a B_L(x) df(x). \quad (8.19)$$

Letting $a \rightarrow -\infty$, this yields

$$0 \leq \lim_{a \rightarrow -\infty} |g(a)| [f(a) - f(-\infty)] \leq \lim_{a \rightarrow -\infty} \int_{-\infty}^a B_L(x) df(x) = 0 \quad (8.20)$$

and

$$\lim_{a \rightarrow -\infty} g(a) [f(a) - f(-\infty)] = 0. \quad (8.21)$$

(b) The existence of the limit $\lim_{x \rightarrow \infty} f(x)$ entails that the integral $\int_b^\infty df(x) = f(\infty) - f(b)$ also exists. Since $f(x)$ is monotonic nondecreasing on the interval (M, ∞) and $\int_M^\infty B_U(x) df(x)$ exists, we get from (5.25): for $b > M$,

$$0 \leq \int_b^\infty |g(b)| df(x) \leq \int_b^\infty B_U(x) df(x) \quad (8.22)$$

hence

$$0 \leq |g(b)| [f(\infty) - f(b)] \leq \int_b^\infty B_U(x) df(x). \quad (8.23)$$

Letting $b \rightarrow \infty$, this yields

$$0 \leq \lim_{b \rightarrow \infty} |g(b)| [f(\infty) - f(b)] \leq \lim_{b \rightarrow \infty} \int_b^\infty B_U(x) df(x) = 0 \quad (8.24)$$

and

$$\lim_{b \rightarrow \infty} g(b) [f(\infty) - f(b)] = 0. \quad (8.25)$$

□

7.1. See von Bahr and Esseen (1965), Chung (1974, p. 48) and Chow and Teicher (1988, p. 108).

7.2. See Chung (1974, p. 48).

PROOF OF PROPOSITION 7.2 The first inequality follows by recursion on applying the Minkowski inequality for two variables. The first part of the second inequality is obtained by multiplying both sides of the first one by $(1/n)$. The second part follows on observing that the function $x^{1/r}$ is concave in x for $x > 0$ when $r > 1$. □

7.3. See von Bahr and Esseen (1965, Theorem 1).

7.4. See von Bahr and Esseen (1965, Theorem 2).

7.5. See von Bahr and Esseen (1965, Theorem 3).

7.6. See von Bahr and Esseen (1965, Theorem 4).

References

- BEEBACK, P. R. (1984): "Inequalities for Absolute Moments of a Distribution: From Laplace to von Mises," *Journal of Mathematical Analysis and Applications*, 98, 435–457.
- CHOW, Y. S., AND H. TEICHER (1988): *Probability Theory. Independence, Interchangeability, Martingales. Second Edition*. Springer-Verlag, New York.
- CHUNG, K. L. (1974): *A Course in Probability Theory*. Academic Press, New York, second edn.
- DEVINATZ, A. (1968): *Advanced Calculus*. Holt, Rinehart and Winston, New York.
- FELLER, W. (1966): *An Introduction to Probability Theory and its Applications, Volume II*. John Wiley & Sons, New York.
- HAASER, N. B., AND J. A. SULLIVAN (1991): *Real Analysis*. Dover Publications, New York.
- HANNAN, E. J. (1985): "Letter to the Editor: "Sufficient and Necessary Conditions for Finite Negative Moments"," *The American Statistician*, 39, 326.
- HONG, L. (2012): "A Remark on the Alternative Expectation Formula," *The American Statistician*, 66, 232–233.
- (2015): "Another Remark on the Alternative Expectation Formula," *The American Statistician*, 69, 157–159.
- KHURI, A., AND G. CASELLA (2002): "The Existence of the First Negative Moment Revisited," *The American Statistician*, 56, 44–47.
- LEHMANN, E. L., AND J. P. SHAFFER (1988): "Inverted Distributions," *The American Statistician*, 42, 191–194.
- LO, A. (2019): "Demystifying the Integrated Tail Probability Formula," *The American Statistician*, 73(4), 360–366.
- LOÈVE, M. (1977): *Probability Theory, Volumes I and II*. Springer-Verlag, New York, 4th edn.
- PIEGORSCH, W. W., AND G. CASELLA (1988): "The Existence of the First Negative Moment," *The American Statistician*, 39, 60–62.
- PROTTER, M. H., AND C. B. MORREY (1991): *A First Course in Real Analysis*, Undergraduate Texts in Mathematics. Springer-Verlag, New York, second edn.
- RIESZ, F., AND B. SZ.-NAGY (1955/1990): *Functional Analysis*. Dover Publications, New York, second edn.
- SERFLING, R. J. (1980): *Approximation Theorems of Mathematical Statistics*. John Wiley & Sons, New York.
- VON BAHR, B., AND C.-G. ESSEEN (1965): "Inequalities for the r th Absolute Moment of a Sum of Random Variables, $1 \leq r \leq 2$," *The Annals of Mathematical Statistics*, 36, 299–303.